

E 6392



00006392



Reg. No.....

Name.....

B.Sc. DEGREE (C.B.C.S.S.) EXAMINATION, MAY 2024

Fourth Semester

Complementary Course—Physics

PHYSICAL OPTICS, LASER PHYSICS AND ASTROPHYSICS

(Common for B.Sc. Mathematics (Model I) and B.Sc. Statistics)

[2013 to 2016 Admissions]

Time : Three Hours

Maximum : 60 Marks

Part A

Answer all questions.

Each question carries 1 mark.

1. Why are interference fringes formed by Newton's rings are circular ?
2. What are the conditions for producing observable interference pattern ?
3. What is the optical path difference between the waves emerging from the slits in the Young's double experiment ?
4. What is double refraction ?
5. State the law of Malus.
6. What are the applications of polaroids ?
7. What are Einstein coefficients ?
8. What are the application of laser beams ?

(8 × 1 = 8)

Part B

Answer any six questions.

Each question carries 2 marks.

9. Explain why different colours are produced by a thin film in white light.
10. Differentiate between diffraction and interference.
11. What is dispersive power ? Explain.
12. What are quarter wave plates ? What are its uses ?

Turn over





E 6392

13. How would you obtain plane polarised light by reflection ?
14. Explain the principle of ruby laser.
15. Give a brief note on Neutron star.
16. Explain population inversion.
17. Discuss about polarization by scattering.
18. Write a short note on the properties of laser beams.

(6 × 2 = 12)

Part C

*Answer any **four** questions.
Each question carries 4 marks.*

19. Two straight and narrow parallel slits 1 mm apart are illuminated by monochromatic light. Fringes formed on the screen held at a distance of 100 cm from the slits are 0.50 mm apart. What is the wavelength of light?
20. In Young's double slit experiment, the angular width of a fringe formed on a screen is 0.1° . The wavelength of light used is 6000 \AA . What is the spacing between the slits?
21. Newton's rings are formed with red light of wavelength 670nm . The radius of the 20th ring is found to be $1.1 \times 10^{-2} \text{ m}$. Find the radius of curvature of the 30th ring.
22. In Fraunhofer diffraction due to a narrow slit a screen is placed 2 m away from the lens to obtain the pattern. If the slit width is 0.2 mm and the first minima lie 5 mm on either side of the central maxima, find the wavelength of light.
23. In a plane diffraction grating, the angle of diffraction for the second order maxima for wavelength $5 \times 10^{-5} \text{ cm}$ is 30° . Calculate the number of lines per cm of the grating surface.
24. The He-Ne systems capable of lasing at 3.391 micrometre. Determine the energy difference in electron volt between the upper and lower levels of the wavelength.

(4 × 4 = 16)

Part D

*Answer any **two** questions.
Each question carries 12 marks.*

25. Discuss the theory of Newton's rings and determination of wavelength.
26. Distinguish between plane polarized, circularly polarized and elliptically polarized light.
27. Explain the principles, construction and working of a YAG laser.
28. Write an essay on stellar evolution.

(2 × 12 = 24)

